

An Articulatory Analysis of American English /ɪ/ in Typically Developing Children and Children Receiving Ultrasound Biofeedback

Amanda Eads¹, Heather Kabakoff¹, Hannah King², Jon Preston³, & Tara McAllister¹
New York University, Université Paris Cité, Syracuse University
November 3, 2022
Ultrafest X


Background

American English /ɹ/

- Articulation:
 - Literature has typically discussed /ɹ/ in terms of a binary “bunched” or “retroflex” shape.

(Espy Wilson & Boyce, 1999;
Zhou et al., 2008;
Archangeli et al., 2011;
Mielke et al., 2016)

**More
Bunched**




**More
Retroflex**

FIGURE 1. The Delattre and Freeman /ɹ/ taxonomy (Delattre & Freeman 1968:41), printed with the permission of Walter de Gruyter GmbH. Copyright 2009.

Figure: Mielke et al. (2016)

American English /ɪ/

- Tongue shapes for /ɪ/ vary within as well as across speakers with a greater number of bunched only speakers in American English (Mielke et al., 2016).



American English /ɪ/

- Typically developing children exhibit broadly adult-like /ɪ/ patterns, but there is large variation between individuals and across /ɪ/ contexts (McGowan et al., 2004; Magloughlin, 2016).





Figure: Two male twins producing a bunched and retroflex shape for “pear” Magloughlin (2016)

American English /ɪ/

- Receives a lot of attention in the clinical literature because it is later developing and considered one of the most difficult sounds to master.
- Ball et al. (2013) describes retroflexion as easier and more commonly cued during treatment.
- Flipsen (2021) describes a “Speech Buddies” device commonly used to aid /ɪ/ development. It cues retroflex tongue shapes.



Ultrasound Biofeedback

- A systematic review demonstrated positive effects for using ultrasound biofeedback in treatment of residual speech sound disorder for /ɪ/ (RSSD; Sugden et al., 2019).
- So, what is it?
 - Client views their tongue on the screen during portions of treatment.
 - Clinician initially cues bunched and retroflex to see which receives better response.
 - Template is laid on screen for client to match during productions.



Research Questions

- Do children show within- and between-speaker variation in tongue shape patterns for /ɹ/ consistent with expectations from adult studies?
- Do patterns differ across groups of children with and without history of RSSD affecting /ɹ/?

Methods

Group 1: With History of RSSD

- 30 American English speakers aged 9-15 who received treatment for /ɪ/
 - 17 males and 13 females
 - Additional 4 participants were excluded due to zero productions of perceptually accurate /ɪ/
- Three phases of 10-week treatment study:
 - 1 week of intensive motor-based treatment (3x)
 - 1 week of intensive ultrasound biofeedback treatment (3x)
 - 8 weeks of low-intensity ultrasound biofeedback treatment (2x per week)

Group 2: Without History of RSSD

- 36 American English speakers aged 9-15 who had no known speech, language, or hearing issues
 - 17 males and 19 females
- Completed the same evaluation tasks as the biofeedback group in two 2-hour sessions on separate days
- Ultrasound data collected with 3-D printed helmet (Derrick et al., 2018)

Syllable Repetition Task

- 45 opportunities for /ɪ/ production (Miccio 2002):
 - Syllabic and postvocalic /ɪ/ in 9 syllables/disyllables

mɪ	dɪ	ʒg / ʒ	ɪr	ɛr	aɪə	ɔr	ar	auə
----	----	--------	----	----	-----	----	----	-----

- Prevocalic /ɪ/ in 6 CV syllables

ri	re	raɪ	ru	ro	ɾa
----	----	-----	----	----	----

Syllable Repetition Task

- During the task, the clinician provided a visual and auditory model of each syllable and cued the child to repeat with their “best /ɪ/ sound”; the clinician also scored each production.
- All targets were repeated 3 times each
- All targets differed from the wordlists used in all other tasks

Articulatory Coding

- Get Contours and SLURP in the ultrasound tracing processes (Tiede, 2020; LaPorte & Ménard, 2018).
- Images selected were the most representative /ɪ/ images nearest to the temporal midpoint of the acoustically measured steady state.

Articulatory Coding




Fig. 3. Decision tree used to classify tongue shapes into five distinct categories for /r/ from ultrasound data.

Figure: /ɹ/ classification tree from King and Ferragne (2020) which includes two bunched shapes, two retroflex shapes, and one in-between (front-up).

Articulatory Coding





Fig. 2. Raw ultrasound frames showing typical examples of each of the five /r/ configurations. The tongue tip is on the right side of the image. The top two images are bunched, while the bottom three are retroflex. The final retroflex configuration exhibits curling up of the tongue with a bright white line where the tongue tip is expected.

King and Ferragne (2020)

Articulatory Coding

- Developed a PsychoPy Coding Program following King & Ferragne 2020
- Viewed concatenated target image + 2 preceding and following images while coding
- Images did not include tongue tracing
- All concatenated images from both groups were randomized for blind classification



Results

General Coding Results

- Total of 2698 coded images
 - 1078 images for the group with history of RSSD
 - 1620 images for the group without history of RSSD
- Bunched vs. Retroflex Ratio
 - 65.2% bunched (1760)
 - 29% retroflex (783)
 - 5.7% flagged (155)


RQ1: Do children show within- and between-speaker variation in tongue shape patterns for /ɹ/ consistent with expectations in adult studies?

■ Patterns appear consistent!

- Adults n= 27 (Mielke et al., 2016)
- Children n =66


■ Tongue Shape types were grouped into 3 categories:

- Bunched Only
- Mixed – Used both types
- Retroflex Only



RQ2: Do patterns differ across groups of children with and without history of RSSD affecting /ɹ/?


- Patterns are not consistent when children are split into groups.
- Number of Participants in each group:
 - Without History of RSSD = 36
 - With History of RSSD = 30
 - Adults = 27



RQ2: Do patterns differ across groups of children with and without history of RSSD affecting /ɹ/?

■ Group Differences


- Statistically significant ($\beta = -10.99$, $SE = 1.91$, $p < .001$) according to mixed effects logistic regression: $\text{Bunched_or_Retroflex} \sim \text{Group} + (1|\text{Subject})$



Discussion

Why the significant difference in tongue shapes?

- Could this be an effect of clinician cueing?
 - Clinicians cued both bunched and retroflex shapes at the beginning of treatment.



Why the significant difference in tongue shapes?

- Are retroflex tongue shapes easier to acquire in the context of treatment for RSSD?
 - Note: our data only include the perceptually accurate tokens.
 - Follow up analysis on positional and vowel influence.



Why the significant difference in tongue shapes?

- Could this be related to articulatory patterning for /ɹ/ in non-rhotic dialects?
 - Children with RSSD are functionally similar to speakers of non-rhotic dialects of English.

- Higher rates of retroflexion have been found in non- rhotic English dialects in New Zealand and the UK (Heyne et al., 2018; King & Ferragne, 2020).
 - Retroflex is favored in the prevocalic contexts.
 - Bunched is favored in the postvocalic contexts.
 - These contexts are “r- less” in non-rhotic dialects
 - Therefore, non-rhotic speakers experience less articulatory pressure to bunch than their rhotic counterparts.

In Conclusion

- This finding furthers our understanding of tongue shape types for /ɪ/ among American English-speaking children 9-15 years old.
 - Very few studies have explored /ɪ/ tongue shapes in children – usually in much younger ages.
 - None have compared across children who learned to produce perceptually accurate /ɪ/ during treatment and typically developing children.

- This study has the potential to inform clinical practice concerning retroflex versus bunched tongue shape cueing.

Thank you!!

- To you for listening and feedback!
- Many thanks to my collaborators, committee members, and BITS lab assistants for help with data processing.
- Special thanks to Chris Carignan and Jeff Mielke for PCA+LDA advice.

Questions?

Please feel free to contact me: are326@nyu.edu

References

- Archangeli, D., Baker, A., & Mielke, J. (2011). *Categorization and features*. In R. Ridouane (Ed.), *Where do phonological features come from? : Cognitive, physical and developmental bases of distinctive speech categories* (pp. 175-195). John Benjamins Publishing Company.
- Ball, M. J., Muller, N., & Granese, A. (2013). Towards an evidence base for /r/-therapy in English. *Journal of Clinical Speech and Language Studies*, 20, 1–23.
- Boyce, S. (2015). The Articulatory Phonetics of /r/ for Residual Speech Errors. *Seminars in Speech and Language*, 36(04), 257–270. <https://doi.org/10.1055/s-0035-1562909>
- Campbell, H., & McAllister Byun, T. (2018). Deriving individualised /r/ targets from the acoustics of children's non-rhotic vowels. *Clinical Linguistics & Phonetics*, 32(1), 70–87. <https://doi.org/10.1080/02699206.2017.1330898>
- D. Derrick, C. Carignan, W.-r. Chen, M. Shujau, C.T. Best Three-dimensional printable ultrasound transducer stabilization system *J. Acoust. Soc. Am.*, 144 (5) (2018), pp. EL392-EL398
- Espy-Wilson, C., & Boyce, S. (1999). A simple tube model for American English /r/. *ICPhS-14*, 2137-2140.
- Flipsen Jr., P. (2021). *Remediation of /r/ for Speech Language Pathologists*. Plural Publishing, Inc.
- Heyne, M., Wang, X., Derrick, D., Dorreen, K., & Watson, K. (2018). The articulation of /ɹ/ in New Zealand English. *Journal of the International Phonetic Association*, 1–23. <https://doi.org/10.1017/S0025100318000324>.
- King, H., & Ferragne, E. (2020). Loose lips and tongue tips: The central role of the /r/-typical labial gesture in Anglo-English. *Journal of Phonetics*, 80, 100978. <https://doi.org/10.1016/j.wocn.2020.100978>
- Magloughlin, L. (2016). Accounting for variability in North American English /ɹ/: Evidence from children's articulation. *Journal of Phonetics*, 54, 51–67. <https://doi.org/10.1016/j.wocn.2015.07.007>
- McGowan, R. S., Nittrouer, S., & Manning, C. J. (2004). Development of [ɹ] in young, Midwestern, American children. *The Journal of the Acoustical Society of America*, 115(2), 871–884. <https://doi.org/10.1121/1.1642624>
- Miccio, A. W. (2002). Clinical Problem Solving: Assessment of Phonological Disorders. *American Journal of Speech-Language Pathology*, 11(3), 221–229.
- Mielke, J., Baker, A., & Archangeli, D. (2016). Individual-level contact limits phonological complexity: Evidence from bunched and retroflex /ɹ/. *Language*, 92(1), 101–140. <https://doi.org/10.1353/lan.2016.0019>
- Laporte, C., & Ménard, L. (2018). Multi-hypothesis tracking of the tongue surface in ultrasound video recordings of normal and impaired speech. *Medical image analysis*, 44, 98–114.
- Peirce, J. W., Gray, J. R., Simpson, S., MacAskill, M. R., Höchenberger, R., Sogo, H., Kastman, E., Lindeløv, J. (2019). PsychoPy2: experiments in behavior made easy. *Behavior Research Methods*. 10.3758/s13428-018-01193-y
- Sugden, E., Lloyd, S., Lam, J., & Cleland, J. (2019). Systematic review of ultrasound visual biofeedback in intervention for speech sound disorders. *International Journal of Language & Communication Disorders*, 54(5), 705–728. <https://doi.org/10.1111/1460-6984.12478>
- Tiede, M. (2020). GetContours: tongue contour fitting software; v3.3. [Computer program]. <https://github.com/mktiede/GetContours>.
- Zhou, X., Espy-Wilson, C. Y., Boyce, S., Tiede, M., Holland, C., & Choe, A. (2008). A magnetic resonance imaging-based articulatory and acoustic study of “retroflex” and “bunched” American English /r/. *J. Acoust. Soc. Am.*, 123(6), 16